



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Reply

Papageorgiou, Spyridon N ; Koretsi, Vasiliki ; Jäger, Andreas

DOI: <https://doi.org/10.1093/ejo/cjw066>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-133162>

Journal Article

Accepted Version

Originally published at:

Papageorgiou, Spyridon N; Koretsi, Vasiliki; Jäger, Andreas (2017). Reply. European Journal of Orthodontics, 39(1):107.

DOI: <https://doi.org/10.1093/ejo/cjw066>

Title Page

Reply

Spyridon N. Papageorgiou^{1,2}, Vasiliki Koretsi³, Andreas Jäger⁴

¹Clinic of Orthodontics and Pediatric Dentistry, Center of Dental Medicine, University of Zurich, Zurich, Switzerland

²Department of Oral Technology, School of Dentistry, University of Bonn, Germany

³Department of Orthodontics, School of Dentistry, University Medical Centre Regensburg, Germany

⁴Department of Orthodontics, School of Dentistry, University of Bonn, Germany

Correspondence to: Spyridon N. Papageorgiou, Clinic of Orthodontics and Pediatric Dentistry, Center of Dental Medicine, Plattenstrasse 11, 8032, University of Zurich, Zurich, Switzerland, E-mail: snpapage@gmail.com

Words in text: 3993

Acknowledgements

SNP is supported by a research grant from Forschungsgemeinschaft Dental (FGD) e.V.

Conflicts of interest

None declared.

Manuscript

Dear Sir,

We thank Dr. Saltaji for the interest in our empirical study on the role of historical control groups in orthodontics (1) and for the opportunity to explain our findings, focusing on the main comparison of our study (i.e. historical versus concurrent controls).

First, the concern expressed by Dr. Saltaji is understandable, but not necessarily wholly justified. By careful consideration of the forest plot for the main analysis (Figure 2), it is evident that there is a large variation in effect magnitude of the included meta-analyses, but not in the effect direction of the included meta-analyses (i.e. all black boxes lie on different gray contours, but are always to the left of the central line). Additionally, no statistical heterogeneity can be measured ($I^2=0\%$). Therefore, if this wide array of different interventions and outcomes indicates consistency, this might be expected to contribute to the credibility results. Of course this might be negated in the future by additional meta-analyses with completely opposite effects, but this has always been the case of evolution of evidence in the scientific procedure. However, based on existing evidence, the 95% predictive intervals (2) indicate that we might expect the Δ SMD for historical controls in future meta-analysis to lie between -0.56 and -0.07 (Table 2); therefore it would still be consistent.

Second, Dr. Saltaji correctly identifies the importance of appropriate basic study design, as this might influence both the magnitude (3) and statistical significance (4) of their results. However, this has been taken into account in our study through the use of multivariable regression methods to adjust for meta-confounders (5). To put it plainly, we calculated the influence of the control group on the study results, while at the same time controlling for the confounding effect of the basic study design (i.e. whether the study was prospective and retrospective). The use of this technique enabled calculation of more precise estimates (6); though with the cost of having fewer eligible meta-analyses.

Third, we comprehend the appeal of setting a minimum number of trials per meta-analysis. It must be however made clear, that the number of 5 trials / meta-analysis that is suggested is an arbitrarily set number and that neither the cited Egger et al. (7) study nor the original commentary of Clarke (8), upon which the statement of the former is based, provide any empirical evidence for adopting such a cut-off in this or future meta-epidemiological studies.

Fourth, as originally reported, *post hoc* power analysis with the method recently proposed by Girardeau et al. (9) indicated that 4-7 extra studies would improve statistical power (although this technique does not precisely reflect the multivariable nature of our analysis plan). However, contrary to Dr. Saltaji's statement, there is no evidence whatsoever that the results of our meta-epidemiological study are heterogeneous, as I^2 was 0% and all studies were consistent on their direction.

The plea of Dr. Saltaji for more meta-analyses including clinical trials of interventions with historical control groups to inform future empirical assessments might make sense from an epistemological point of view, but based on existing evidence could be clinically misleading. Given the state of our evidence base (10), every attempt should be made to improve the quality of orthodontic meta-analyses (11) and to safeguard our clinical recommendations from known *or suspected* sources of bias.

References

1. Papageorgiou, S.N., Koretsi, V. and Jäger, A. (2016) Bias from historical control groups used in orthodontic research: a meta-epidemiological study. *European Journal of Orthodontics*. First published online on April 29, 2016, 10.1093/ejo/cjw035.
2. Int'Hout, J., Ioannidis, J.P., Rovers, M.M. and Goeman, J.J. (2016) Plea for routinely presenting prediction intervals in meta-analysis. *BMJ Open*, 6, e010247.
3. Papageorgiou, S.N., Xavier, G.M. and Cobourne, M.T. (2015) Basic study design influences the results of orthodontic clinical investigations. *Journal of Clinical Epidemiology*, 68, 1512–1522.
4. Papageorgiou, S.N., Kloukos, D., Petridis, H. and Pandis, N. (2015) Publication of statistically significant research findings in prosthodontics & implant dentistry in the context of other dental specialties. *Journal of Dentistry*, 43, 1195–202.
5. Siersma, V., Als-Nielsen, B., Chen, W., Hilden, J., Gluud, L.L. and Gluud, C. (2007) Multivariable modelling for meta-epidemiological assessment of the association between trial quality and treatment effects estimated in randomized clinical trials. *Statistics in Medicine*, 26, 2745–58.
6. Kelley, K. and Maxwell, S.E. (2003) Sample size for multiple regression: obtaining regression coefficients that are accurate, not simply significant. *Psychological Methods*, 8, 305–21.
7. Egger, M., Juni, P., Bartlett, C., Holenstein, F. and Sterne, J. (2003) How important are comprehensive literature searches and the assessment of trial quality in systematic reviews? Empirical study. *Health Technology Assessment*, 7, 1–76.
8. Clarke, M. (2002) Searching for trials for systematic reviews: what difference does it make? *International Journal of Epidemiology*, 31, 123–4.
9. Giraudeau, B., Higgins, J.P., Tavernier, E. and Trinquart, L. (2016) Sample size calculation for meta-epidemiological studies. *Statistics in Medicine*, 35, 239–250.
10. Papageorgiou, S.N., Papadopoulos, M.A. and Athanasiou, A.E. (2014) Reporting characteristics of meta-analyses in orthodontics: methodological assessment and statistical recommendations. *European Journal of Orthodontics*, 36, 74–85.
11. Papageorgiou, S.N. (2014) Meta-analysis for orthodontists: Part II—Is all that glitters gold? *Journal of Orthodontics*, 41, 327–336.